


2024 Adult Drug Reference

Drug	Indication	Dosing	Cautions	Comments
Acetaminophen	Pain Control	IV/IO – 1,000 mg over 15 minutes (1 drop/sec w/ 10 drop/ml tubing – administer over 15 min)	Do not administer to patients with severe liver impairment/ disease.	Do not administer if patient has taken more than 2,000 mg in past 24 hours.
Adenosine	Narrow complex tachycardia	Initial – 6 mg rapid IV Repeat – 12 mg rapid IV Follow each dose with 20 ml NS rapid IV	May cause transient heart block or asystole. Use ½ dose for patients taking carbamazepine or dipyridole. Do not administer if patient is experiencing acute asthma exacerbation.	Side effects include: chest pressure/pain, palpitations, hypotension, dyspnea, or feeling of impending doom.
Albuterol	Bronchospasm	5 mg nebulized or 6 puffs MDI with spacer Repeat as needed	Use continuous cardiac monitoring with patients taking MAOI antidepressants (Phenelzine and Tarnylcypromine)	Use patient’s MDI, if available. When using MDI, always use a spacer.
	Crush injury/Hyperkalemia	10 mg nebulized continuously or 6 puffs MDI with spacer		
Aspirin	Chest pain – suspected cardiac or STEMI	324 mg PO	Contraindicated in aspirin or salicylate allergy	Blood thinner use is not a contraindication.
Atropine	Symptomatic bradycardia	Initial – 1 mg IV/IO Repeat every 3-5 min. to a max of 3 mg	Doses less than 0.5mg can cause paradoxical bradycardia.	Can dilate pupils, aggravate glaucoma, cause urinary retention, confusion, and dysrhythmias including V-Tach and V-Fib. Increases myocardial oxygen consumption. Bradycardia in children is primarily related to respiratory issues – assure adequate ventilation first.
	Organophosphate overdose	Initial – 1-2 mg IV/IO/IM Repeat every 3-5 min. until relief of symptoms is achieved		
Calcium Chloride	Calcium channel blocker OD	1 g IV/IO over 60 seconds	Use cautiously or not at all in patients on digitalis. Avoid extravasation. Rapid administration can cause dysrhythmias or arrest.	None
	Crush injury			
	Hyperkalemia			
Dextrose 10%	Hypoglycemia	Initial – 100 ml IV Repeat – 150 ml if glucose remains ≤ 70 mg/dl and patient remains altered	Can cause tissue necrosis if IV is infiltrated	Recheck blood glucose after administration.
Diphenhydramine	Allergic reaction	50 mg IV/IO/IM	None	May cause drowsiness.
	Nausea in pregnancy < 20 weeks	25 mg IV/IO/IM		
	Dystonic reaction	25-50 mg IV/IO or 50 mg IM		
Dopamine	Persistent hypotension unrelated to hypovolemia	400 mg in 250 ml NS Infuse at 5-20 mcg/kg/min titrated to response	None	None
Epi 1:1,000	Anaphylaxis	0.3 mg IM May repeat x 1 after 5 min.	Never administer IV/IO. Use with caution in asthma patients with a history of hypertension or coronary artery disease. May cause serious dysrhythmias or exacerbate angina.	None
	Asthma/COPD or respiratory distress			
	Stridor	5 mg nebulized		
Epi 1:10,000	PEA/Asystole V-Fib/Pulseless V-Tach	1 mg IV/IO May repeat every 3-5 min. to a max of 3 mg	May cause serious dysrhythmias or exacerbate angina. In adult anaphylactic patients, should be used if patient is hypotensive or no improvement after two (2) Epi 1:1,000 IM doses.	None
	Anaphylaxis (bradycardia after max Epi 1:1,000 IM)	0.1 mg slow IV/IO May repeat every 3-5 min. as needed to a max of 0.5 mg		
EpiPen	Allergic reaction/Anaphylaxis	1 auto-injector	None	None



2024 Adult Drug Reference

Drug	Indication	Dosing	Cautions	Comments
Fentanyl	Pain control	IV/IO – 50 mcg initial May repeat every 5 min. to max of 200 mcg or IM – 50 mcg initial May repeat x1 after 10 min. to max of 100 mcg or IN – 50 mcg (½ each nare) Do not repeat	Contraindicated in SBP < 90mmHg, child birth or active labor, sudden onset of severe headache, AMS, suspected closed head injury. Can cause hypotension or respiratory depression. If initial dose is via IN, may repeat by other route to maximum indicated – must account for initial dose when calculating maximum dose.	Recheck vital signs between each dose. Hypotension is more common in patients with low cardiac output or volume depletion. Respiratory depression is reversible with naloxone.
Glucagon	Hypoglycemia	IM – 1 mg May repeat x 1 after 10 min.	None	Effect may be delayed 15-20 minutes.
	Symptomatic Beta Blocker overdose	IV/IO/IM – 1-3 mg Do not repeat		
Glucose paste	Hypoglycemia	24 g PO	Not indicated with AMS or if patient cannot swallow.	None
Glucola	Hypoglycemia	50 g PO	Not indicated with AMS or if patient cannot swallow.	None
Levalbuterol	Bronchospasm	2.5 mg nebulized Repeat as needed	Use continuous cardiac monitoring with patients taking MAOI antidepressants (Phenelzine and Tarnylcypromine).	None
	Crush injury/Hyperkalemia	5 mg nebulized continuously		
Lidocaine	Persistent V-Fib/V-Tach	1-1.5 mg/kg IV/IO May repeat 0.5-0.75 mg/kg to a maximum of 3 mg/kg	Use caution with bradycardia. Can cause cardiac dysrhythmia	None
	IO anesthetic	40 mg IO		
Midazolam	Seizure	IM – 10 mg May repeat x1 after 5 min. or IN – 10 mg (½ each nare) May repeat x1 after 5 min. or IV if already established – 5 mg May repeat x1 after 5 min.	Use caution in patients over 60 years of age.	Monitor respiratory status after administration.
	Hyperactive delirium	5 mg IM/IN May repeat x1 after 5 min.		
	Sedation for pacing or cardioversion	2.5 mg IV/IO May repeat to a max of 5 mg		
	Sedation of patient with an advanced airway	2.5 mg IV/IO May repeat to a max of 5 mg		
	Dizziness/vertigo	2.5 mg IV/IN	Do not administer if patient is > 50 for dizziness/vertigo.	
Naloxone	Respiratory depression or apnea associated with suspected opioid overdose	IN – 2 mg ½ dose each nare or IM/IV - 1-2 mg	Abrupt withdrawal symptoms and combative behavior may occur.	IN administration preferred unless patient is in shock or has copious secretions/blood in nares. Titrate to effect of normal respirations; it is not necessary to fully wake the patient.
Naloxone autoinjector/preload	Overdose	1 preload syringe	See Naloxone	See Naloxone
Nitroglycerin	Chest pain	0.4 mg SL May repeat as needed every 5 min.	Can cause hypotension and headache. Do not administer if systolic BP < 110mmHg or heart rate < 50. Do not administer if patient has taken Viagra, Levitra, Staxyn, or Stendra within past 24 hours or Cialis if taken within 36 past hours.	None
	Pulmonary edema	0.4 mg SL if systolic BP > 110mmHg 0.8 mg SL if systolic BP > 150mmHg May repeat appropriate dose every 5 min.		
Ondansetron	Vomiting or severe nausea	4 mg IV/IO/IM/ODT May repeat after 15 min. to a maximum of 12 mg	Administer IV/IO dose over 1 minute as rapid administration may cause syncope.	For patients with nausea who are < 20 weeks pregnant, consider Diphenhydramine.
	Nausea in pregnancy > 20 weeks			
Sodium Bicarbonate	Tricyclic antidepressant overdose	1 mEq/kg IV/IO	Can precipitate with or inactivate other drugs.	Use only if life-threatening or in the presence of hemodynamically significant dysrhythmias.
	Crush injury			
	Hyperkalemia			
	Cardiac arrest with known renal failure			
Suboxone	Opioid withdrawal	Base Hospital order required SL – 16 mg (sublingual) May repeat x 1 with 8 mg after 10 min. Maximum dose = 24 mg	None	If patient requests to AMA after administration, recontact Base Hospital.
Valium	Hazmat/WMD exposure	Refer to dosing guide attached to ChemPak kit	None	None

