

Seizure – Post

For any seizure that stopped prior to EMS arrival and there is no further seizure activity during EMS contact

History


- Reported or witnessed seizure
- Previous seizure history
- Medical alert tag
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse, or abrupt cessation
- Fever

Signs and Symptoms

- Altered mental status
- Sleepiness
- Incontinence
- Evidence of trauma
- Unconscious
- Incontinence
- Bitten tongue/oral trauma

Differential

- Head trauma
- Metabolic, hepatic or renal failure
- Tumor
- Hypoxia
- Electrolyte abnormality
- Drugs or medication non-compliance
- Infection or sepsis
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia
- Epilepsy
- Syncope



Pearls

- Status Epilepticus is defined as two or more successive seizures without a period of consciousness or recovery, or one prolonged seizure lasting longer than 5 minutes. This is a true emergency requiring rapid airway control, treatment, and transport.
- Assess the possibility of occult trauma and substance abuse.
- Be prepared for airway problems and continued seizures.
- Be prepared to assist ventilations or manage the airway, especially if Midazolam is used.

