

Pediatric Stroke/CVA/TIA

For suspected stroke or transient ischemic attack (stroke symptoms that resolve rapidly)

History


- Last seen normal
- A&O Status and GCS
- Family members phone number
- Previous stroke or TIA or brain hemorrhage
- Major surgery within last 2 weeks
- Signs of active bleeding, including Melena
- Associated diseases (DM, HTN, CAD)
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- History of brain tumor, aneurysm, or AVM.

Signs and Symptoms

- Altered mental status
- Weakness or paralysis
- Blindness or other sensory loss
- Aphasia or dysarthria
- Syncope
- Vertigo or dizziness
- Vomiting
- Headache
- Seizure
- Respiratory pattern change
- Hypertension/hypotension
- Diplopia or double vision

Differential

- See Altered Mental Status
- TIA
- Sepsis
- Seizure/Todd's paralysis
- Hypoglycemia
- Stroke
 - Thrombotic or embolic (~85%)
 - Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis or renal failure
- Bell's Palsy



Primary Stroke Centers
 Kaiser South San Francisco
 Sequoia Hospital
 Mills-Peninsula
 Kaiser Redwood City
 Stanford Hospital

Thrombectomy Capable Stroke Centers
 Mills-Peninsula

Comprehensive Stroke Centers
 Kaiser Redwood City
 Stanford Hospital

Pediatric Stroke/CVA/TIA

For suspected stroke or transient ischemic attack (stroke symptoms that resolve rapidly)

Pearls

- Pediatric strokes do occur.
- Time last known well: One of the most important items that prehospital providers can obtain, on which all treatment decisions are based. Be very precise in gathering data to establish the time of onset and report as an actual time (i.e., "13:45," NOT "about 45 minutes ago"). Without this information, patients may not be able to receive thrombolytics at the hospital. For patients who "woke up and noticed stroke symptoms," time starts when the patient was last awake.
- The differential listed on the Altered Mental Status protocol should also be considered.
- Be alert for airway problems (difficulty swallowing, vomiting and aspiration). PO meds are not appropriate.
- Hypoglycemia or hyperglycemia can present as a LOCALIZED neurologic deficit.
- Document the Cincinnati Prehospital Stroke Scale in the EHR.

