

Pediatric Cardiac Arrest

For non-traumatic cardiac arrest in which any resuscitation is initiated, NOT dead on arrival

History


- Code status (DNR or POLST)
- Events leading to arrest
- Estimated downtime
- History of current illness
- Past medical history
- Medications
- Existence of terminal illness

Signs and Symptoms

- Unresponsive
- Apneic
- Pulseless

Differential

- Airway obstruction/respiratory disease
- Medical vs. trauma
- VF vs. pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory arrest or drug overdose



Pediatric Cardiac Arrest Treatment Protocols

Pediatric Cardiac Arrest

For non-traumatic cardiac arrest in which any resuscitation is initiated, NOT dead on arrival

Pearls

- Airway is a more important intervention in pediatric arrests. This should be accomplished quickly with a BVM, airway adjunct, and appropriately sized mask. Patient survival is often dependent on proper ventilation and oxygenation.
- Efforts should be directed at high quality chest compressions with limited interruptions.
- Use appropriately sized pediatric BVM with EtCO₂.
- Do not delay chest compressions while applying any device or intervention.
- Use a metronome during chest compression to ensure proper rate.
- Provide resuscitative efforts for 30 minutes to maximize chance of ROSC.
- If resuscitative efforts do not attain ROSC, consider cessation of efforts per Operations 10 – Determination of Death.
- Resuscitation is based on proper planning and organized execution. Procedures require space and patient access. Make room to work. Utilize a team focused approach assigning responders to predetermined tasks.
- Reassess airway and document EtCO₂ frequently.
- Defibrillation vests should be removed by EMS personnel before compressions, but do not cut vests. Once removed, disengage battery to prevent alarming.
- Pediatric pads should be used in children < 10kg or measurement of Purple.

