

# Pediatric V-Fib/Pulseless V-Tach

For non-traumatic cardiac arrest in which any resuscitation is initiated, NOT dead on arrival

**History**

- Events leading to arrest
- Estimated downtime
- Prior resuscitation attempts
- Past medical history
- Medications
- Known terminal illness

**Signs and Symptoms**

- Pulseless
- Apneic

**Differential**

- Airway obstruction/respiratory disease
- Medical vs. trauma
- VF vs. pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory arrest or drug overdose

 Enter from Cardiac Arrest

P	<b>Defibrillation</b>
	Use <i>Broselow Tape</i> ; refer to dosing guide Resume chest compressions ( <b>15:2 ratio</b> ) 1.5 inches for infants; 2 inches for children Change compressors every 2 minutes (Limit changes/pulse checks to < 5 seconds)
	Establish IV/IO

**AT ANY TIME**  
Return of spontaneous circulation  
  
Go to Post Resuscitation

P	<b>Defibrillation</b>
	Use <i>measuring tape</i> ; refer to dosing guide Resume chest compressions ( <b>15:2 ratio</b> ) 1.5 inches for infants; 2 inches for children Change compressors every 2 minutes (Limit changes/pulse checks to < 5 seconds)
	<b>Epinephrine (1:10,000)</b> Use <i>Broselow Tape</i> ; refer to dosing guide

P	<b>Defibrillation</b>
	Use <i>measuring tape</i> ; refer to dosing guide Resume chest compressions ( <b>15:2 ratio</b> ) 1.5 inches for infants; 2 inches for children Change compressors every 2 minutes (Limit changes/pulse checks to < 5 seconds)
	<b>Lidocaine</b> Use <i>Broselow Tape</i> ; refer to dosing guide

Persistent V-Fib/V-Tach

 Aystole/PEA

Return of spontaneous circulation?

 Post Resuscitation

**Notify receiving facility.  
Consider Base Hospital for  
medical direction**

Pediatric Cardiac Arrest Treatment Protocols



# Pediatric V-Fib/Pulseless V-Tach

For non-traumatic cardiac arrest in which any resuscitation is initiated, NOT dead on arrival

## Pearls

- Airway is a more important intervention in pediatric arrests. This should be accomplished quickly with a BVM, airway adjunct, and appropriately sized mask. Patient survival is often dependent on proper ventilation and oxygenation.
- Efforts should be directed at high quality chest compressions with limited interruptions.
- Use appropriately sized pediatric BVM with EtCO<sub>2</sub>.
- Do not delay chest compressions while applying any device or intervention.
- Use a metronome during chest compression to ensure proper rate.
- Provide resuscitative efforts for 30 minutes to maximize chance of ROSC.
- If resuscitative efforts do not attain ROSC, consider cessation of efforts per Operations 10 – Determination of Death.
- Resuscitation is based on proper planning and organized execution. Procedures require space and patient access. Make room to work. Utilize a team focused approach assigning responders to predetermined tasks.
- Reassess airway and document EtCO<sub>2</sub> frequently.
- Defibrillation vests should be removed by EMS personnel before compressions, but do not cut vests. Once removed, disengage battery to prevent alarming.

