

Pediatric Carbon Monoxide/Cyanide

For suspected or known carbon monoxide exposure

History

- Industrial or closed space fire
- Facial burns
- Previous CO poisoning
- Propane powered equipment (e.g., power mower, tractor, gas powered equipment)
- Gas home heaters, natural gas stoves, kerosene heaters
- Gas clothes dryer or hot water heater
- Multiple people or pets with similar symptoms

Signs and Symptoms



- AMS
- Malaise/Fatigue
- Flu-like symptoms
- Weakness
- Headache
- Dizziness
- Blurred vision
- Ataxia
- Seizure
- Nausea/vomiting/cramping
- Chest pain

Differential

- Diabetic emergency
- Infection/sepsis
- Myocardial infarction
- Anaphylaxis
- Renal failure
- Head injury/trauma
- Ingestion/toxic exposure

E	Immediately remove from exposure
	Airway support
	High flow Oxygen regardless of SpO ₂
P	Blood glucose analysis
	Cardiac monitor
	CO-oximetry (SpCO), if available
	12-Lead ECG
	<i>Consider, IV</i>
	Consider EtCO ₂ monitoring
<i>Consider, Normal Saline bolus</i> <i>Use length-based tape; refer to dosing guide</i> May repeat x2	

[Emergency Hyperbaric Chambers](#)
 John Muir Medical Center – Walnut Creek

 **Notify receiving facility.**
Consider Base Hospital for medical direction 

Pearls

- For suspected cyanide poisoning, contact the receiving hospital early.
- CO is colorless and odorless.
- Pulse oximetry will likely be normal with CO toxicity.
- Hyperbaric oxygen is recommended for those with AMS, seizure, coma, focal deficits, blindness, CO levels > 25% or > 20% if pregnant. John Muir Medical Center – Walnut Creek is the only emergency hyperbaric chamber in the Bay Area. Contact the Base Hospital for direction.
- Consider cyanide poisoning in any patient with CO intoxication.
- Consider cyanide poisoning in any patient with AMS.