

## **SNAKEBITE**

APPROVED: Gregory Gilbert, MD EMS Medical Director  
Nancy Lapolla EMS Director

DATE: July 2018

### **Information Needed:**

- Type of snake, if known and location found
- Appearance of snake, shape of pupil, presence of stripes or rattle, size of snake
- Time of bite
- Prior first aid by patient or friends
- Symptoms: local pain or swelling, metallic taste in mouth, hypotension, coma, bleeding

### **Objective Findings:**

- One or more punctures wounds, or horseshoe set of teeth marks
- For pit-vipers (*Crotalines*), there is a spectrum of envenomation from non-envenomation to serious envenomation

### **Non-envenomated:**

- No discoloration around puncture marks
- Little or no pain after a few minutes

#### **Treatment:**

- Safety first; do not attempt to capture snake and do not handle an apparently dead snake or decapitated snake head with your hands
- If transporting the snake, be certain that it is in a closed solid container
- Remove rings or other jewelry which might constrict circulation later
- Routine Medical Care
- Transport all suspected patients with envenomations for medical evaluations

### **Serious Envenomation:**

- Dark discoloration around punctures within 5 minutes
- Marked edema formation
- Severe pain within a short time
- Altered mental status
- Oozing of hemolyzed blood from punctures, possible formation of fluid blebs on skin
- Fasciculation
- Hypotension
- Marked tachycardia
- Definite metallic taste

Treatment:

- Safety first; do not attempt to capture snake
- Remove rings or other jewelry which might constrict circulation later
- Routine Medical Care
- Monitor EKG
- Transport all suspected envenomations for medical evaluations
- Document distal pulse
- Immobilize bitten part with splint, etc.
- Consider IV access in the unaffected extremity
- Fluid challenge 250-1000 ml NS for hypotension
- Consider pain management-see Interim Adult and Pediatric Pain Assessment and Management protocol (June 2018).

**Precautions and Comments:**

- Do not incise envenomations
- If the snake is dead, bring it in for positive identification in a closed solid container. Avoid the fangs because they are capable of envenomation even when dead. If alive, do not try to capture
- Ice applied directly to skin surfaces can cause serious tissue damage and should not be used
- All so-called “pet” snakes should be identified
- Exotic poisonous snakes such as those in zoos have different signs and symptoms than those of the pit vipers. Zoos and legal exotic snake collectors are required to have a starter supply of antivenom on hand for each type of snake. Bring the antivenom with the patient to the hospital
- Bites from coral snakes, *elapids* related to cobras, usually do not have any early symptoms, thus all bites are considered envenomated (coral snakes are not indigenous to California)
- Early notification of the receiving hospital is recommended