

Multi-System Trauma

For any traumatic injuries that involve multiple systems or isolated chest or abdominal injuries. For injuries involving the head, use Head Trauma protocol

History


- Time of injury
- Mechanism (blunt vs. penetrating)
- Damage to structure or vehicle
- Location of patient in structure or vehicle
- Restraints or protective equipment use
- Past medical history
- Medications

Signs and Symptoms

- Evidence of trauma
- Pain, swelling, deformity, lesions, or bleeding
- AMS
- Unconscious
- Respiratory distress or failure
- Hypotension or shock
- Arrest

Differential

- Chest:
 - Tension pneumothorax
 - Flail chest
 - Pericardial tamponade
 - Open chest wound
 - Hemothorax
- Intra-abdominal bleeding
- Pelvis or femur fracture
- Spinal injury
- Head injury
- Hypothermia



Adult and Pediatric Trauma Treatment Protocols

Multi-System Trauma

For any traumatic injuries that involve multiple systems or isolated chest or abdominal injuries. For injuries involving the head, use Head Trauma protocol

Pearls

- ALS procedures in the field do not significantly improve patient outcome in critical trauma patients.
- Basic airway management is preferred unless unable to effectively manage with BLS maneuvers. Utilize modified jaw thrust technique to open the airway.
- Intubation of head injury patients is best addressed at the hospital.
- Hypotension is age dependent and is not always a reliable sign. It should be interpreted in context with the patient's typical BP, if known. Shock may be present with a seemingly normal blood pressure initially.
 - Neonate: < 60mmHg or weak pulses
 - Infant: < 70mmHg or weak pulses
 - 1-10 years: < 70mmHg + (age in years x2)
 - Over 10 years: <80mmHg
 - Over 65 years: <110mmHg
- Stabilize flail segments with bulky dressing.
- Cover eviscerated bowel with dry sterile dressing.
- Stabilize impaled object(s) with bulky dressing. Do not remove.
- Avoid hyperventilation. Maintain an EtCO₂ of 35 or greater, which may be unreliable if the patient was subject to multisystem trauma or poor perfusion.
- An important item to monitor and document is a change in the level of consciousness by repeat examination.
- Do not overlook the possibility of associated domestic violence or abuse.

