


Traumatic Arrest


For cardiac arrest with penetrating or blunt traumatic mechanism. NOT for trauma sustained after cardiac arrest, use primary impression Cardiac Arrest – Non-traumatic

History <ul style="list-style-type: none"> Evidence of trauma or blood loss Events leading to arrest Estimated downtime 	Signs and Symptoms <ul style="list-style-type: none"> Unresponsive Apneic Pulseless 	Differential <ul style="list-style-type: none"> Tension pneumothorax Cardiac tamponade Hypovolemic shock Spinal shock Traumatic brain injury
---	---	--

E <p>Has a paramedic, EMT, designated first responder, or public safety officer found injuries incompatible with life, including one or more of the following conditions:</p> <ul style="list-style-type: none"> Decapitation Incineration Rigor Mortis Decomposition Apnea with destruction and/or separation of the body from the heart, brain, liver, or lungs Multi-casualty incidents (MCIs) where triage principles preclude the initiation or continuation of resuscitation 	<p style="text-align: center;">AT ANY TIME</p> <p style="text-align: center;">Return of spontaneous circulation</p>  <p style="text-align: center;">Multi-System Trauma</p>
---	---

!

Tourniquet use should not be delayed until a patient is in shock or is clearly exsanguinating. It should be applied early and can be used safely without risk of patient injury. Do not wait; apply often and tighten if needed.



Adult and Pediatric Trauma Treatment Protocols

Pearls

- Patients who do not qualify for field determination of death but have or develop cardiopulmonary arrest should be transported to the closest trauma center.